

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 2

In the Claims

Please amend the claims by replacing all prior versions of the claims pursuant to 37 C.F.R. §1.121 as modified by 68 Fed. Reg. 38611 (June 30, 2003) as indicated below.

1. (Currently Amended) A method of modifying the content or composition, or content and composition, of a metabolite in the ~~storage organ seed~~ of a plant, ~~said metabolite selected from the group consisting of fatty acid, starch, soluble non-starch polysaccharide, insoluble non-starch polysaccharide, fibre, and total protein nitrogen, said method comprising:~~

- (i) expressing in the ~~storage organ seed~~ of the plant a chimeric gene comprising a nucleotide sequence encoding a sulfur-rich protein placed operably in connection with a promoter sequence capable of conferring expression in ~~said storage organ the seed~~, wherein the sulfur-rich protein is a 2S protein or the Aspl synthetic protein;
- (ii) determining the content or composition, or content and composition, of a metabolite in ~~said storage organ the seed~~, said metabolite selected from the group consisting of fatty acid, starch, soluble non-starch polysaccharide, insoluble non-starch polysaccharide, fibre and total protein nitrogen, and
- (iii) selecting a plant having a modified content or composition, or content and composition, of ~~said a~~ metabolite in the ~~storage organ seed~~ thereof, as compared to a plant in which said chimeric gene is not expressed,  
wherein the plant is wheat, oats, maize, barley,

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 3

rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea, and

wherein the modified content or composition, or content and composition, of a metabolite in the seed of a plant is at least one of:

- a) increased total protein nitrogen content;
- b) decreased total starch content;
- c) increased or decreased fatty acid content;
- d) modified fatty acid composition;
- e) increased or decreased fibre content; or
- f) modified fibre composition.

2-4. (Canceled)

5. (Previously presented) The method according to claim 1 wherein the total protein nitrogen content is increased.

6. (Previously presented) The method according to claim 1 wherein the fibre content or composition is modified.

7. (Canceled)

8. (Previously presented) The method according to claim 1 wherein the fatty acid content is increased or decreased.

9-10. (Canceled)

11. (Previously presented) The method according to claim 1 wherein the sulfur-rich protein is Brazil Nut Protein (BNP).

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 4

12. (Previously presented) The method according to claim 1 wherein the sulfur-rich protein is sunflower seed albumin (SSA).
13. (Previously presented) The method according to claim 1 wherein the sulfur-rich protein is the Aspl synthetic protein.
14. (Previously presented) The method according to claim 1 wherein the plant is a dicotyledonous plant.
15. (Previously presented) The method according to claim 14 wherein the dicotyledonous plant is a pea or chickpea plant.
16. (Previously presented) The method according to claim 1 wherein the promoter sequence comprises the pea vicilin gene promoter sequence.
17. (Previously presented) The method according to claim 1 wherein the plant is a monocotyledonous plant.
18. (Original) The method according to claim 17 wherein the monocotyledonous plant is a rice plant.
19. (Previously presented) The method according to claim 1 wherein the promoter comprises a *Triticum aestivum* HMW glutenin promoter sequence.

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 5

20. (Previously presented) The method according to claim 1 further comprising the first steps of:

- (i) introducing the chimeric gene into a plant cell, tissue, organ or whole organism; and
- (ii) regenerating an intact plant therefrom.

21. (Currently Amended) A method of increasing the total protein nitrogen content of seeds of a plant, said method comprising

- (i) expressing in the seeds of the plant a chimeric gene comprising a nucleotide sequence encoding a sulfur-rich protein placed operably in connection with a promoter sequence capable of conferring expression in said seeds, wherein the sulfur-rich protein is a 2S protein or the Aspl synthetic protein, said nucleotide sequence also positioned upstream of a transcription termination sequence;

(ii) determining the ~~level~~ content of total protein nitrogen in the seeds; and

~~(iii)~~ (iii) selecting a plant having an increased total protein nitrogen content in the seeds thereof as compared to the seeds of a plant which does not express the chimeric gene,

wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

22. (Original) The method according to claim 21 wherein the promoter sequence is the pea vicilin gene promoter and the plant is a dicotyledonous plant.

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 6

23. (Original) The method according to claim 21 wherein the chimeric gene further comprises the pea vicilin gene promoter and transcription terminator sequences.
24. (Previously presented) The method according to claim 22 wherein the plant is pea or chickpea.
25. (Original) The method according to claim 21 wherein the promoter sequence is a wheat HMW glutenin gene promoter and the plant is a monocotyledonous plant.
26. (Original) The method according to claim 21 wherein the chimeric gene further comprises the wheat HMW glutenin gene promoter and/or NOS transcription terminator sequences.
27. (Previously presented) The method according to claim 25 wherein the plant is a rice plant.
28. (Currently Amended) A method of ~~modifying the~~ increasing or decreasing fatty acid content of seeds of a plant, said method comprising:
- (i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds:
  - (ii) determining the ~~level~~ content of a fatty acid in the seed; and

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 7

(iii) selecting a plant having ~~a modified increased or decreased content of the fatty acid content~~ in the seeds thereof as compared to the seeds of a plant which does not express the chimeric gene, wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

29. (Original) The method according to claim 28 wherein the promoter sequence is the pea vicilin gene promoter and the plant is a dicotyledonous plant.

30. (Original) The method according to claim 28 wherein the chimeric gene further comprises the pea vicilin gene promoter and transcription terminator sequences.

31. (Canceled)

32. (Previously presented) The method according to claim 29 wherein if the plant is pea the ~~level~~ content of fatty acids in the seeds is decreased.

33. (Currently Amended) A method of modifying the fatty acid composition of seeds of a plant, said method comprising  
(i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 8

- gene in the seeds of said plant;
- (ii) determining the fatty acid composition of the seeds;  
and
- (iii) selecting a plant having a modified fatty acid composition in the seeds thereof, as compared to seeds of a plant in which the chimeric gene is not expressed.

wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

34. (Original) The method according to claim 33 wherein the promoter sequence is the pea vicilin gene promoter and the plant is a dicotyledonous plant.
35. (Original) The method according to claim 33 wherein the chimeric gene further comprises the pea vicilin gene promoter and transcription terminator sequences.
36. (Canceled)
37. (Previously presented) The method according to claim 33 wherein the fatty acid is selected from the group consisting of: myristic acid, stearic acid, gadoleic acid, behenic acid, lignoceric acid, oleic acid, linoleic acid, linolenic acid and erucic acid.
38. (Currently Amended) A method of decreasing the starch content of seeds of a plant, said method comprising:
- (i) expressing in the seeds of said plant a chimeric

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 9

gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant:

- (ii) determining the starch content of the seeds; and
- (iii) selecting a plant having a decreased starch content in the seeds thereof, as compared with seeds of a plant in which the chimeric gene is not expressed, wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

- 39. (Original) The method according to claim 38 wherein the promoter sequence is the pea vicilin gene promoter and the plant is a dicotyledonous plant.
- 40. (Original) The method according to claim 38 wherein the chimeric gene further comprises the pea vicilin gene promoter and transcription terminator sequences.
- 41. (Previously presented) The method according to claim 39 wherein the plant is a pea plant.
- 42. (Currently Amended) A method of modifying the amino acid composition of seeds of a plant, said method comprising:
  - (i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed



Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 10

upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant;  
(ii) determining the amino acid composition of the seeds; and

(iii) selecting a plant having a modified amino acid composition in the seeds thereof, as compared to a plant in which the chimeric gene is not expressed,

wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

43. (Original) The method according to claim 42 wherein the promoter sequence is the pea vicilin gene promoter and the plant is a dicotyledonous plant.
44. (Original) The method according to claim 42 wherein the chimeric gene further comprises the pea vicilin gene promoter and transcription terminator sequences.
45. (Previously presented) The method according to claim 43 wherein the plant is pea or chickpea.
46. (Original) The method according to claim 45 wherein the proportion of arginine relative to other amino acids is increased.

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 11

47. (Currently Amended) A method of ~~modifying~~ increasing or decreasing the fibre content of seeds of a plant, said method comprising:

- (i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant;
- (ii) determining the fibre content of the seeds; and
- (iii) selecting a plant having ~~a modified~~ an increased or a decreased content of fibre content in the seeds thereof, as compared with seeds of a plant which does not express said chimeric gene,  
wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

48. (Original) The method according to claim 47 wherein the promoter sequence is the pea vicilin gene promoter and the plant is a dicotyledonous plant.

49. (Original) The method according to claim 47 wherein the chimeric gene further comprises the pea vicilin gene promoter and transcription terminator sequences.

50. (Canceled)

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 12

51. (Previously presented) The method according to claim 48 wherein if the plant is a pea plant the ~~level~~ content of fibre in the seed is increased.

52. (Currently Amended) A method of modifying the fibre quality composition of seeds of a plant, said method comprising:

- (i) the step of expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant;
- (ii) determining the fibre quality composition of the seeds; and
- (iii) selecting a plant having a modified fibre quality composition in the seeds thereof, as compared with seeds of a plant in which the chimeric gene is not expressed.

wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

53. (Original) The method according to claim 52 wherein the promoter sequence is the pea vicilin gene promoter and the plant is a dicotyledonous plant.

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 13

54. (Original) The method according to claim 53 wherein the chimeric gene further comprises the pea vicilin gene promoter and transcription terminator sequences.

55. (Canceled)

56. (Currently Amended) The method according to claim 52 wherein the ~~level~~ content of soluble ~~NSP and/or non-starch polysaccharide, or the level content~~ of insoluble ~~NSP non-starch polysaccharide, or the content of both~~ in the seed is decreased.

57. (Currently Amended) The method according to claim 52 wherein the ~~level~~ content of lignin in the seed is increased.

58-63. (Canceled)

64. (Currently Amended) The method according to any one of claims ~~1, 21, 28, 33, 38, 42, 47, or 52~~ further comprising the first steps of:

- (i) introducing the chimeric gene into a plant cell, tissue, organ or whole organism; and
- (ii) regenerating an intact plant therefrom.

65. (Currently Amended) A transformed plant ~~produced having a modified content or composition, or content and composition, of a metabolite in its seed, wherein the modification is~~ by the method according to any one of claims 1, 21, 28, 33, 38, 42, 47, 52, or 102, or progeny of said plant, wherein said progeny comprises at least

Applicants: Higgins et al.  
Serial No.: 09/508,579  
Filed : May 10, 2000  
Page : 14

one copy of the chimeric gene in an expressible format ~~7~~  
~~said plant being wheat, oats, maize, barley, rice,~~  
~~sorghum, millet, oilseed rape, rye, safflower, sunflower,~~  
~~potato, sweet potato, beetroot, taro, Jerusalem~~  
~~artichoke, onion, garlic, soybean, pea, bean (including~~  
~~joba and common bean), lentil, chickpea or sugar cane.~~

66. (Canceled)

67. (Previously presented) A ~~plant part~~ seed derived from the  
plant according to claim 65 wherein said ~~plant part~~ seed  
comprises at least one copy of the chimeric gene ~~present~~  
~~in said plant or progeny~~ in an expressible format.

68. (Canceled)

69. (Original) The ~~plant part according to claim 67~~  
~~consisting of seeds~~ seed derived from the transformed  
plant of claim 65, wherein the seed has a modified  
content or composition, or content and composition of a  
metabolite, wherein the modification is at least one of:

- a) increased total protein nitrogen content;
- b) decreased total starch content;
- c) increased or decreased fatty acid content;
- d) modified fatty acid composition;
- e) increased or decreased fibre content; or
- f) modified fibre composition.

70-87. (Canceled)

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 15

88. (Currently Amended) The method of claim 1 wherein the chimeric gene encodes sunflower seed albumin (SSA) and ~~wherein the storage organ is a seed~~ and wherein the total protein nitrogen content of the seed is increased.
89. (Currently Amended) The method of claim 1 wherein the chimeric gene encodes sunflower seed albumin (SSA) and ~~wherein the storage organ is a seed~~ and wherein the amino acid composition of the seed is modified.
90. (Currently Amended) The method of claim 1 wherein the chimeric gene encodes sunflower seed albumin (SSA) and ~~wherein the storage organ is a seed~~ and wherein expression of the chimeric gene in the seed increases or decreases the fibre content of the seed.
91. (Currently Amended) The method of claim 1 or claim 102 wherein the chimeric gene encodes sunflower seed albumin (SSA) and ~~wherein the storage organ is a seed~~ and wherein expression of the chimeric gene in the seed modifies the fibre composition of the seed.
92. (Currently Amended) The method of claim 1 wherein the chimeric gene encodes sunflower seed albumin (SSA) and ~~wherein the storage organ is a seed~~ and wherein expression of the chimeric gene in the seed decreases the total starch content of the seed.

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 16

93. (Currently Amended) The method of claim 1 wherein the chimeric gene encodes sunflower seed albumin (SSA) and ~~wherein the storage organ is a seed~~ and wherein expression of the chimeric gene in the seed increases or decreases the total fatty acid content of the seed.
94. (Currently Amended) The method of claim 1 wherein the chimeric gene encodes sunflower seed albumin (SSA) and ~~wherein the storage organ is a seed~~ and wherein expression of the chimeric gene in the seed modifies the fatty acid composition of the seed.
95. (Canceled)
96. (Previously presented) The method of claim 1 wherein the chimeric gene encodes sunflower seed albumin (SSA) and further comprises a promoter sequence which confers strong expression at least in the seeds of the plant.
97. (Previously presented) The method according to claim 96 wherein the promoter is the pea vicilin promoter.
98. (Previously presented) The method according to claim 96 wherein the promoter is the wheat HMW glutenin promoter.
99. (Previously presented) The method of claim 96 wherein the chimeric gene further comprises a transcription terminator sequence placed downstream of the sequence encoding SSA.

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 17

100. (Previously presented) The method of claim 99 wherein the transcription terminator sequence is the pea vicilin gene terminator sequence.

101. (Currently Amended) The method according to claim 1 or claim 102, wherein the content or composition, or content and composition, of more than one metabolite in the ~~storage organ seed~~ of the plant is modified, ~~and wherein at least one of said metabolites is selected from the group consisting of fatty acid, starch, soluble non starch polysaccharide, insoluble non starch polysaccharide, fibre and total protein nitrogen.~~

102. (Currently Amended) A method of modifying the content or composition, or content and composition, of a metabolite in the ~~storage organ seed~~ of a plant, ~~said metabolite selected from the group consisting of fatty acid, starch, soluble non starch polysaccharide, insoluble non starch polysaccharide, fibre, and total protein nitrogen, said method comprising:~~

- (i) expressing in the ~~storage organ seed~~ of the plant a chimeric gene comprising a nucleotide sequence encoding a sulfur-rich protein placed operably in connection with a promoter sequence capable of conferring expression in ~~said storage organ the seed~~, wherein the sulfur-rich protein is a 2S protein or the Aspl synthetic protein; and
- (ii) determining whether the content or composition, or content and composition, of a metabolite in ~~said storage organ the seed of the plant is modified as compared to that of a plant in which the chimeric gene is not expressed~~, ~~said metabolite selected~~



Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 18

~~from the group consisting of fatty acid, starch, soluble non-starch polysaccharide, insoluble non-starch polysaccharide, fibre and total protein nitrogen~~

wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea, and

wherein the modified content or composition, or content and composition, of a metabolite in the seed of a plant is at least one of:

- a) increased total protein nitrogen content;
- b) decreased total starch content;
- c) increased or decreased fatty acid content;
- d) modified fatty acid composition;
- e) increasing or decreasing fibre content; or
- f) modified fibre composition.

103. (Currently Amended) A method of increasing the total protein nitrogen content of seeds of a plant, said method comprising:

- (i) expressing in the seeds of the plant a chimeric gene comprising a nucleotide sequence encoding a sulfur-rich protein placed operably in connection with a promoter sequence capable of conferring expression in said seeds, wherein the sulfur rich protein is a 2S protein or the Aspl synthetic protein, said nucleotide sequence also positioned upstream of a transcription termination sequence; and
- (ii) determining whether the level content of total protein nitrogen in the seeds of said plants the

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 19

plant is increased as compared to that of a plant in which the chimeric gene is not expressed,  
wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

104. (Currently Amended) A method of modifying increasing or decreasing the fatty acid content of seeds of a plant, said method comprising:

- (i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds; and
- (ii) determining the level whether the content of a fatty acid in the seeds of the plant is increased or decreased as compared to that of a plant in which the chimeric gene is not expressed,  
wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

105. (Currently Amended) A method of modifying the fatty acid composition of seeds of a plant, said method comprising:

- (i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 20

upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant; and

- (ii) determining whether the fatty acid composition of the seeds of the plant is modified as compared to that of a plant in which the chimeric gene is not expressed,

wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

106. (Currently Amended) A method of decreasing the starch content of seeds of a plant, said method comprising:

- (i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant; and
- (ii) determining whether the starch content of the seeds of the plant is decreased as compared to that of a plant in which the chimeric gene is not expressed,
- wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 21

107. (Currently Amended) A method of modifying the amino acid composition of seeds of a plant, said method comprising:

- (i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant; and
- (ii) determining whether the amino acid composition of the seeds of the plant is modified as compared to that of a plant in which the chimeric gene is not expressed.

wherein the plant is wheat, oats, maize, barley, rice, sorghum, millet, rye, safflower, sunflower, soybean, pea, bean, faba bean, common bean, lentil, or chickpea.

108. (Currently Amended) A method of modifying increasing or decreasing the fibre content of seeds of a plant, said method comprising:

- (i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant; and
- (ii) determining whether the fibre content of the seeds of the plant is increased or decreased as compared to that of a plant in which the chimeric gene is not

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 22

expressed,

wherein the plant is wheat, oats, maize, barley,  
rice, sorghum, millet, rye, safflower, sunflower,  
soybean, pea, bean, faba bean, common bean, lentil, or  
chickpea.

109. (Currently Amended) A method of modifying the fibre quality composition of seeds of a plant, said method comprising:

(i) expressing in the seeds of said plant a chimeric gene that comprises a structural gene sequence encoding sunflower seed albumin (SSA) placed upstream of a transcription termination sequence and operably in connection with a promoter sequence capable of conferring expression on said structural gene in the seeds of said plant; and

(ii) determining whether the fibre quality composition of the seeds of the plant is modified as compared to that of a plant in which the chimeric gene is not expressed,

wherein the plant is wheat, oats, maize, barley,  
rice, sorghum, millet, rye, safflower, sunflower,  
soybean, pea, bean, faba bean, common bean, lentil, or  
chickpea.

110. (Currently Amended) The method according to claim 1, wherein the total protein nitrogen content of ~~storage organ~~ the seed of the plant is increased by at least 10%.

111. (Canceled)

Applicants: Higgins et al.  
Serial No.: 09/508,979  
Filed : May 10, 2000  
Page : 23

112. (Currently Amended) The method according to claim 1, wherein the total fibre content of the ~~storage organ~~ seed of the plant is increased or decreased by at least 5%.

113. (Currently Amended) The method according to claim 1, wherein the total starch content of ~~storage organ~~ the seed of the plant is reduced by at least 10%.

114. (Currently Amended) The method according to claim 1, wherein the total fatty acid content of ~~storage organ~~ the seed of the plant is increased or decreased by at least 5%.

115. (Currently Amended) The method according to claim 1, wherein the content of any one fatty acid in the ~~storage organ~~ seed of the plant is increased or decreased by at least 5%.